

MATH 128 = Calculus 2 for the Sciences, Fall 2006
Assignment 1

Due September 20 at the beginning of class
postponed to Fri. Sept. 22/06

Evaluate the following integrals **using the method indicated**.

Note: on a test or exam you will not typically be given the method of integration to use.

Justify each answer, showing all necessary steps. Simplify your answers as much as possible (without a calculator, of course).

1. Use u -substitution (also referred to as "change of variable"):

(a) $\int \frac{(\ln x)^2}{x} dx$

(b) $\int_0^1 x^3 e^{1-x^4} dx$

(c) $\int_0^{\frac{\pi}{2}} \frac{\sin t}{1+\cos t} dt$

2. Use integration by parts:

(a) $\int e^{-\theta} \sin 3\theta d\theta$

(b) $\int_0^1 s \tan^{-1} s ds$

3. Use a suitable trigonometric substitution:

(a) $\int_0^{2\sqrt{3}} \frac{x^3}{\sqrt{16-x^2}} dx$

(b) $\int \frac{dt}{(1+t^2)^{3/2}}$

4. Use partial fractions:

(a) $\int \frac{10x+20}{(x^2-1)(x^2+4)} dx$

(b) $\int_1^4 \frac{2p^2+3}{p(p+1)^2} dp$

①

$$1. a) I = \int \frac{(\ln x)^2}{x} dx \quad u = \ln x \Rightarrow du = \frac{dx}{x}$$

$$\Rightarrow I = \int u^2 du = \frac{u^3}{3} + C = \frac{(\ln x)^3}{3} + C$$

$$b) I = \int_0^1 x^3 e^{-x^4} dx$$

$$\text{Solution 1: } I = e \int_0^1 x^3 e^{-x^4} dx$$

$$\text{Let } u = -x^4 \Rightarrow du = -4x^3 dx \Rightarrow -\frac{du}{4} = x^3 dx,$$

$$\left[x=0 \Rightarrow u=0, x=1 \Rightarrow u=-1 \right] \quad \text{②}$$

$$\Rightarrow I = e \int_0^{-1} \left(-\frac{du}{4} \right) e^u = -\frac{e}{4} \int_0^{-1} e^u du \quad \text{①}$$

Note the reversal

of integration limits requires multiplying the integral by -1 .

(see page 36b text.)

$$= \frac{e}{4} \int_{-1}^0 e^u du = \frac{e}{4} [e^u]_{-1}^0$$

$$= \frac{e}{4} (e^0 - e^{-1}) = \frac{e}{4} (1 - e^{-1})$$

$$= \frac{e}{4} - \frac{1}{4} \quad \text{or} \quad \frac{1}{4} (e - 1)$$

$$\text{or Solution 2: } I = \int_0^1 x^3 e^{-x^4} dx, \quad u = 1 - x^4, \quad du = -4x^3 dx$$

$$\Rightarrow \int x^3 dx = -\frac{du}{4} \quad \left\{ \begin{array}{l} x=0 \Rightarrow u=1 \\ x=1 \Rightarrow u=0 \end{array} \right. \quad \text{③}$$

$$\Rightarrow I = \int_1^0 \left(-\frac{du}{4} \right) e^u = \frac{1}{4} \int_0^1 e^u du = \frac{1}{4} [e^u]_0^1 = \frac{1}{4} (e - 1) \quad \text{①}$$

$$1c) I = \int_0^{\pi/2} \frac{\sin t \, dt}{1 + \cos t} \quad u = 1 + \cos t \quad \textcircled{1}$$

$$du = -\sin t \, dt$$

$$\text{or } -du = \sin t \, dt \quad \textcircled{1}$$

$$\textcircled{2} \left\{ t=0 \Rightarrow u = 1 + \cos 0 = 1 + 1 = 2, \right.$$

$$\left. t = \frac{\pi}{2} \Rightarrow u = 1 + \cos \frac{\pi}{2} = 1 + 0 = 1 \right\} \quad \textcircled{1}$$

$$\Rightarrow I = \int_2^1 \frac{-du}{u} = \int_1^2 \frac{du}{u} = [\ln u]_1^2$$

$$= \ln 2 - \ln 1 = \ln 2 \quad \textcircled{1}$$

*

$$2a) I = \int e^{-\theta} \sin 3\theta \, d\theta \quad \text{I.B.P.}$$

$$u = \sin 3\theta \quad v = -e^{-\theta} \quad \textcircled{2}$$

$$du = 3\cos 3\theta \, d\theta \quad dv = e^{-\theta} \, d\theta$$

$$\Rightarrow I = -e^{-\theta} \sin 3\theta - \int -e^{-\theta} (3\cos 3\theta) \, d\theta \quad \leftarrow \textcircled{2}$$

$$= -e^{-\theta} \sin 3\theta + 3 \int e^{-\theta} \cos 3\theta \, d\theta \quad \text{I.B.P. again:}$$

$$u = \cos 3\theta \quad v = -e^{-\theta} \quad \textcircled{2}$$

$$du = -3\sin 3\theta \, d\theta \quad dv = e^{-\theta} \, d\theta$$

$$\Rightarrow I = -e^{-\theta} \sin 3\theta + 3 \left[-e^{-\theta} \cos 3\theta - \int -e^{-\theta} (-3\sin 3\theta) \, d\theta \right] \leftarrow \textcircled{2}$$

$$\Rightarrow I = -e^{-\theta} \sin 3\theta - 3e^{-\theta} \cos 3\theta - 9 \int e^{-\theta} \sin 3\theta \, d\theta$$

①

$$\Rightarrow 10I = -e^{-\theta} \sin 3\theta - 3e^{-\theta} \cos 3\theta$$

$$\Rightarrow \boxed{I = -\frac{1}{10} e^{-\theta} \sin 3\theta - \frac{3}{10} e^{-\theta} \cos 3\theta} + C \quad \textcircled{2}$$


The C is ESSENTIAL!

* NOTE: Could let $u = e^{-\theta}$ (both times) instead.

26) $I = \int_0^1 s \tan^{-1} s \, ds$ IBP: $u = \tan^{-1} s$ $v = \frac{s^2}{2}$ (2)

* $du = \frac{ds}{1+s^2}$ $dv = s \, ds$

* Aside



$u = \tan^{-1} s \Rightarrow \tan u = s$

where

$\tan(u(s)) = s \Rightarrow d(\tan u(s)) = d(s)$

$\Rightarrow \frac{d(\sin u(s))}{ds(\cos u(s))} = 1$

Need quotient and chain rule

$\Rightarrow \left(\cos(u(s)) \frac{du}{ds} \right) \cos(u(s)) - \sin(u(s)) \left(\sin(u(s)) \frac{du}{ds} \right) = 1$
 $(\cos^2(u(s))) = 1$

$\Rightarrow \frac{du}{ds} (\cos^2(u(s)) + \sin^2(u(s))) = 1$
 $\cos^2(u(s))$

$\Rightarrow \frac{du}{ds} = \frac{1}{\cos^2(u(s))}$

Triangle $\Rightarrow \cos u(s) = \frac{1}{\sqrt{s^2+1}}$

$\Rightarrow \frac{du}{ds} = \frac{1}{s^2+1} \therefore du = \frac{ds}{s^2+1}$


$\Rightarrow I = \left[\frac{s^2}{2} \tan^{-1} s \right]_0^1 - \int_0^1 \frac{s^2}{2} \frac{ds}{1+s^2}$ (1) (2) (3) LONG DIVISION

$= \left(\frac{1}{2} \tan^{-1} 1 - 0 \right) - \left(\frac{1}{2} \int_0^1 \left(1 - \frac{1}{s^2+1} \right) ds \right)$

$= \frac{1}{2} \left(\frac{\pi}{4} \right) - \frac{1}{2} \left[s - \tan^{-1} s \right]_0^1$ (1)

$= \frac{\pi}{8} - \frac{1}{2} (1 - \tan^{-1} 1 - (0 - \tan^{-1} 0))$ (1)

$= \frac{\pi}{8} - \frac{1}{2} + \frac{1}{2} \left(\frac{\pi}{4} \right) = \left[\frac{\pi}{4} - \frac{1}{2} \right]$ (1)



$$3a) I = \int_0^{2\sqrt{3}} \frac{x^3}{\sqrt{16-x^2}} dx = \int_0^{2\sqrt{3}} \frac{x^3}{\sqrt{4^2-x^2}} dx$$

(14) let $x = 4 \sin \theta$ $\left\{ \begin{array}{l} x=0 \Rightarrow 4 \sin \theta \Rightarrow \theta=0 \\ x=2\sqrt{3} = 4 \sin \theta \Rightarrow \sin \theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = \frac{\pi}{3} \end{array} \right.$

$$\Rightarrow I = \int_0^{\pi/3} \frac{(64 \sin^3 \theta) 4 \cos \theta d\theta}{\sqrt{16-(4 \sin \theta)^2}}$$

$$= \frac{256}{4} \int_0^{\pi/3} \frac{\sin^3 \theta \cos \theta}{\sqrt{16(1-\sin^2 \theta)}} d\theta \quad \leftarrow (1)$$

$$= \frac{256}{4} \int_0^{\pi/3} \frac{\sin^2 \theta \cos \theta}{\cos \theta} d\theta \quad (1)$$

$$= 64 \int_0^{\pi/3} \sin^2 \theta d\theta$$

EXPAND
sin²θ

$$\textcircled{1} = 64 \int_0^{\pi/3} \sin^2 \theta \cos \theta d\theta \quad \leftarrow \cos^2 \theta \sin \theta = 1$$

$$\textcircled{1} = 64 \int_0^{\pi/3} (1 - \cos^2 \theta) \sin \theta d\theta \quad \leftarrow \text{let } u = \cos \theta$$

$du = -\sin \theta d\theta$

$$\textcircled{1} = 64 \int_1^{1/2} -(1-u^2) du$$

$\left\{ \begin{array}{l} \theta=0 \Rightarrow u = \cos 0 = 1 \\ \theta = \frac{\pi}{3} \Rightarrow u = \cos \frac{\pi}{3} = \frac{1}{2} \end{array} \right.$

$$\textcircled{1} = -64 \int_{1/2}^1 (1-u^2) du$$

$$\textcircled{1} = -64 \left[-u + \frac{u^3}{3} \right]_{1/2}^1$$

$$= -64 \left(-1 + \frac{1}{3} - \left(-\frac{1}{2} + \frac{1}{24} \right) \right) = 64 \left(\frac{24-8-12+1}{24} \right)$$


$$\therefore I = \frac{40}{3} \quad (1)$$

3b) $I = \int \frac{dt}{(1+t^2)^{3/2}}$ let $t = \tan \theta$ } (2)
 $dt = \sec^2 \theta d\theta$ } (2)

(9) $\Rightarrow I = \int \frac{\sec^2 \theta d\theta}{(1 + \tan^2 \theta)^{3/2}} = \int \frac{\sec^2 \theta d\theta}{(\sec^2 \theta)^{3/2}}$ (2)

$= \int \frac{\sec^2 \theta d\theta}{\sec^3 \theta} = \int \frac{d\theta}{\sec \theta}$ (1)

$= \int \cos \theta d\theta = \sin \theta + C$ (2)

* Since $t = \tan \theta \Rightarrow$ 

(1) $\Rightarrow \sin \theta = \frac{t}{\sqrt{1+t^2}}$

$\therefore I = \frac{t}{\sqrt{1+t^2}} + C$ (1) **

Note: Since $t = \tan \theta \Rightarrow \theta = \tan^{-1} t$.

Recall: $-\frac{\pi}{2} < \tan^{-1} t < \frac{\pi}{2}$.

So, $\sin \theta = \sin(\tan^{-1} t)$ is defined for all $t \in \mathbb{R}$ (i.e., there are no restrictions on $\sin(\tan^{-1} t)$ that affect our result).

$\therefore I = \sin(\tan^{-1} t) + C$ **

is also correct

(and equivalent to the answer above).

$$4a) I = \int \frac{10x+20}{(x^2-1)(x^2+4)} dx = \int \frac{10x+20}{(x+1)(x-1)(x^2+4)} dx$$

$$= \int \left(\frac{A}{x+1} + \frac{B}{x-1} + \frac{Cx+D}{x^2+4} \right) dx \leftarrow \textcircled{3}$$

$$\textcircled{3} \rightarrow \int \frac{A(x-1)(x^2+4) + B(x+1)(x^2+4) + (Cx+D)(x^2-1)}{(x+1)(x-1)(x^2+4)} dx$$

Since denominators are equal, equate numerators:

$$\textcircled{1} \Rightarrow 10x+20 = A(x-1)(x^2+4) + B(x+1)(x^2+4) + (Cx+D)(x^2-1)$$

$$\Rightarrow 10x+20 = A(x^3 + 4x - x^2 - 4) + B(x^3 + 4x + x^2 + 4) + Cx^3 - (x + Dx^2 - D)$$

$$\Rightarrow 10x+20 = x^3(A+B+C) + x^2(-A+B+D) + x(4A+4B-C) + (-4A+4B-D)$$

Match coefficients on LHS to RHS:

$$x^3: 0 = A+B+C \quad (1)$$

$$x^2: 0 = -A+B+D \quad (2)$$

$$x^1: 10 = 4A+4B-C \quad (3)$$

$$x^0: 20 = -4A+4B-D \quad (4)$$

$$\cdot \text{ Add Eq's (2) + (4)} \Rightarrow 20 = -5A+5B \quad (5)$$

$$\cdot \text{ Add Eq's (1) + (3)} \Rightarrow 10 = 5A+5B \quad (6)$$

$$\cdot \text{ Add Eq's (5) + (6)} \Rightarrow 30 = 10B \Rightarrow B = 3$$

$$\cdot \text{ From Eq (5) (or could use Eq (6))} \Rightarrow 20 = -5A+15 \Rightarrow A = -1$$

$$\cdot \text{ From Eq (1)} \Rightarrow 0 = -1+3+C \Rightarrow C = -2$$

$$\cdot \text{ From Eq (2) (or could use Eq (3))} \Rightarrow 0 = 1+3+D \Rightarrow D = -4$$

- see next page -

7.

4a) cont'd

$$\Rightarrow I = \int \left(\frac{-1}{x+1} + \frac{3}{x-1} + \frac{-2x-4}{x^2+4} \right) dx \quad (1)$$

$$= -\ln|x+1| + 3\ln|x-1| - \int \frac{2x}{x^2+4} + \frac{4}{x^2+4} dx$$

use substitution
 $u = x^2 + 4$, OR, "see

directly" that the
 antiderivative is $\ln(x^2+4)$
 (Since $\frac{d}{dx} \ln(x^2+4) = \frac{2x}{x^2+4}$)

let $x = 2 \tan \theta$
 $\Rightarrow dx = 2 \sec^2 \theta d\theta$
 etc., OR, recall

$$\int \frac{dx}{x^2+a^2} = \int \frac{dx}{a^2 \left(\frac{x}{a} \right)^2 + 1}$$

$$= \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right)$$

So, $\int \frac{4 dx}{x^2+4} = 4 \int \frac{2 \sec^2 \theta d\theta}{4 \tan^2 \theta + 4}$

$$= \frac{8}{4} \int \frac{\sec^2 \theta d\theta}{\tan^2 \theta + 1}$$

$$= 2 \int d\theta = 2\theta + C, \quad \begin{cases} x = 2 \tan \theta \\ \Rightarrow x = \tan \theta \end{cases}$$

$$= 2 \tan^{-1} \left(\frac{x}{2} \right) + C \quad \begin{cases} \Rightarrow \theta = \tan^{-1} \frac{x}{2} \end{cases}$$

$$\therefore I = -\ln|x+1| + 3\ln|x-1| - \ln(x^2+4) + 2 \tan^{-1} \left(\frac{x}{2} \right) + C$$

(5)

4b) $I = \int_1^4 \frac{2p^2+3}{p(p+1)^2} dp$ (3)

(20) $\frac{2p^2+3}{p(p+1)^2} = \frac{A}{p} + \frac{B}{p+1} + \frac{C}{(p+1)^2} = \frac{A(p+1)^2 + Bp(p+1) + Cp}{p(p+1)^2}$ (3)

$\Rightarrow 2p^2+3 = A(p^2+2p+1) + Bp^2 + Bp + Cp \leftarrow (1)$

$\Rightarrow \int p^0: 2 = A+B$ (2) - work
 $\int p^1: 0 = 2A+B+C$ (3) - A, B, C
 $\int p^0: [3=A]$ $\Rightarrow 2 = 3+B \Rightarrow [B = -1]$
 $p^1 \text{ coeff.} \Rightarrow 0 = 2(3) + (-1) + C \Rightarrow [C = -5]$

$\Rightarrow I = \int_1^4 \left(\frac{3}{p} - \frac{1}{p+1} - \frac{5}{(p+1)^2} \right) dp$ (1)

$= [3 \ln|p| - \ln|p+1| + \frac{5}{p+1}]_1^4$ (3)

$= 3 \ln 4 - \ln 5 + \frac{5}{5} - \left(3 \ln 1 - \ln 2 + \frac{5}{2} \right)$

$= 3 \ln 4 - \ln 5 + 1 + \ln 2 - \frac{5}{2}$

$\therefore [I = 3 \ln 4 - \ln 5 + \ln 2 - \frac{3}{2}]$ (4)

OR, since

$\ln 4 = \ln 2^2 = 2 \ln 2,$

$\Rightarrow I = 3(2 \ln 2) - \ln 5 + \ln 2 - \frac{3}{2}$

$= 7 \ln 2 - \ln 5 - \frac{3}{2}$

$= \ln \left(\frac{2^7}{5} \right) - \frac{3}{2}$

} optional.