

## Section 1.1: Systems of Linear Equations

A linear equation:

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$$

**EXAMPLE:**

$$4x_1 - 5x_2 + 2 = x_1 \quad \text{and} \quad x_2 = 2(\sqrt{6} - x_1) + x_3$$

↓

rearranged

↓

$$3x_1 - 5x_2 = -2$$

↓

rearranged

↓

$$2x_1 + x_2 - x_3 = 2\sqrt{6}$$

**Not linear:**

$$4x_1 - 6x_2 = x_1x_2 \quad \text{and} \quad x_2 = 2\sqrt{x_1} - 7$$

A **system of linear equations** (or a **linear system**):

A collection of one or more linear equations involving the same set of variables, say,  $x_1, x_2, \dots, x_n$ .


A **solution** of a linear system:

A list  $(s_1, s_2, \dots, s_n)$  of numbers that makes each equation in the system true when the values  $s_1, s_2, \dots, s_n$  are substituted for  $x_1, x_2, \dots, x_n$ , respectively.

**EXAMPLE** Two equations in two variables:

$$x_1 + x_2 = 10$$


$$-x_1 + x_2 = 0$$



**one unique solution**

$$x_1 - 2x_2 = -3$$


$$2x_1 - 4x_2 = 8$$



**no solution**

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$




**infinitely many solutions**

**BASIC FACT:** A system of linear equations has either


- (i) exactly one solution (*consistent*) or
- (ii) infinitely many solutions (*consistent*) or
- (iii) no solution (*inconsistent*).

**EXAMPLE:** Three equations in three variables. Each equation determines a plane in 3-space.


i) The planes intersect in one point. (*one solution*)



ii) The planes intersect in one line. (*infinitely many solutions*)



iii) There is not point in common to all three planes. (*no solution*)



## The **solution set**:

- The set of all possible solutions of a linear system.

## **Equivalent systems**:

- Two linear systems with the same solution set.

## STRATEGY FOR SOLVING A SYSTEM:

- *Replace one system with an equivalent system that is easier to solve.*

## **EXAMPLE:**

$$x_1 - 2x_2 = -1$$


$$-x_1 + 3x_2 = 3$$

$$x_1 - 2x_2 = -1$$


$$x_2 = 2$$

$$x_1 = 3$$


$$x_2 = 2$$



$$\begin{aligned} x_1 - 2x_2 &= -1 \\ -x_1 + 3x_2 &= 3 \end{aligned}$$



$$\begin{aligned} x_1 - 2x_2 &= -1 \\ x_2 &= 2 \end{aligned}$$



$$\begin{aligned} x_1 &= 3 \\ x_2 &= 2 \end{aligned}$$

## Matrix Notation

$$\begin{array}{rcl} x_1 - 2x_2 & = & -1 \\ -x_1 + 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

(coefficient matrix)

$$\begin{array}{rcl} x_1 - 2x_2 & = & -1 \\ -x_1 + 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 & -1 \\ -1 & 3 & 3 \end{bmatrix}$$

(augmented matrix)

↓

$$\begin{array}{rcl} x_1 - 2x_2 & = & -1 \\ & & x_2 = 2 \end{array} \quad \begin{bmatrix} 1 & -2 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

↓

$$\begin{array}{rcl} x_1 & = & 3 \\ & & x_2 = 2 \end{array} \quad \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \end{bmatrix}$$

## **Elementary Row Operations:**

1. (*Replacement*) Add one row to a multiple of another row.
2. (*Interchange*) Interchange two rows.
3. (*Scaling*) Multiply all entries in a row by a nonzero constant.

**Row equivalent matrices:** Two matrices where one matrix can be transformed into the other matrix by a sequence of elementary row operations.

**Fact about Row Equivalence:** If the augmented matrices of two linear systems are row equivalent, then the two systems have the same solution set.



$$\begin{array}{rcl}
 x_1 & = & 29 \\
 x_2 & = & 16 \\
 x_3 & = & 3
 \end{array}
 \left[ \begin{array}{cccc}
 1 & 0 & 0 & 29 \\
 0 & 1 & 0 & 16 \\
 0 & 0 & 1 & 3
 \end{array} \right]$$

**Solution:** (29, 16, 3)

**Check:** Is (29, 16, 3) a solution of the *original* system?

$$x_1 - 2x_2 + x_3 = 0$$

$$2x_2 - 8x_3 = 8$$

$$-4x_1 + 5x_2 + 9x_3 = -9$$

$$\begin{array}{rcl}
 (29) - 2(16) + 3 & = & 29 - 32 + 3 & = & 0 \\
 2(16) - 8(3) & = & 32 - 24 & = & 8 \\
 -4(29) + 5(16) + 9(3) & = & -116 + 80 + 27 & = & -9
 \end{array}$$

## Two Fundamental Questions (Existence and Uniqueness)

- 1) Is the system consistent; (i.e. does a solution **exist**?)
- 2) If a solution exists, is it **unique**? (i.e. is there one & only one solution?)

**EXAMPLE:** Is this system consistent?

$$\begin{aligned}x_1 - 2x_2 + x_3 &= 0 \\2x_2 - 8x_3 &= 8 \\-4x_1 + 5x_2 + 9x_3 &= -9\end{aligned}$$

In the last example, this system was reduced to the triangular form:

$$\begin{aligned}x_1 - 2x_2 + x_3 &= 0 \\x_2 - 4x_3 &= 4 \\x_3 &= 3\end{aligned} \quad \begin{bmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$

This is sufficient to see that the system is consistent and unique. Why?

**EXAMPLE:** Is this system consistent?

$$\begin{array}{r} 3x_2 - 6x_3 = 8 \\ x_1 - 2x_2 + 3x_3 = -1 \\ 5x_1 - 7x_2 + 9x_3 = 0 \end{array} \quad \left[ \begin{array}{cccc} 0 & 3 & -6 & 8 \\ 1 & -2 & 3 & -1 \\ 5 & -7 & 9 & 0 \end{array} \right]$$

**Solution:** Row operations produce:

$$\left[ \begin{array}{cccc} 0 & 3 & -6 & 8 \\ 1 & -2 & 3 & -1 \\ 5 & -7 & 9 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cccc} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 3 & -6 & 5 \end{array} \right] \rightarrow \left[ \begin{array}{cccc} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 0 & 0 & -3 \end{array} \right]$$

Equation notation of triangular form:

$$\begin{array}{r} x_1 - 2x_2 + 3x_3 = -1 \\ 3x_2 - 6x_3 = 8 \\ 0x_3 = -3 \quad \leftarrow \textit{Never true} \end{array}$$

The original system is inconsistent!

**EXAMPLE:** For what values of  $h$  will the following system be consistent?

$$\begin{aligned}3x_1 - 9x_2 &= 4 \\ -2x_1 + 6x_2 &= h\end{aligned}$$

**Solution:** Reduce to triangular form.

$$\begin{bmatrix} 3 & -9 & 4 \\ -2 & 6 & h \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -3 & \frac{4}{3} \\ -2 & 6 & h \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -3 & \frac{4}{3} \\ 0 & 0 & h + \frac{8}{3} \end{bmatrix}$$

The second equation is  $0x_1 + 0x_2 = h + \frac{8}{3}$ . System is consistent only if  $h + \frac{8}{3} = 0$  or  $h = \frac{-8}{3}$ .